

Foliar Pathogen Growth Response to Elevated CO₂









- Saplings infected:
 - Ambient conditions: 92%
 - Elevated conditions: 83%
- Leaves infected per plant:
 - Ambient: 66%
 - Elevated: 57%

•Leaf area covered by lesions:

-Ambient: 1.66%

-Elevated: 0.78%


Isolating P. minima


Plants produce tannins in greater amounts at higher CO₂ levels and inhibit fungal growth





To see effects of tannic acid on fungal growth:


- •P. minima grown on agar plates in presence of tannic acid
- •Fungal cultures expected to have higher radial growth w/ elevated CO₂
- •TA expected to interfere with fungal growth (but only at high conc?)














×		
	*	
		*significant difference
		S

×	

Conclusions

- Elevated CO₂ reduced infection of red maple saplings by P. minima at the FACE site
 - Reduced # leaves infected
 - Reduced # of lesions on infected leaves
- Although growth area of fungus on agar plates do not appear significantly different in ambient vs. elevated CO₂, trends indicate that a higher tannic acid concentration may inhibit growth over time in elevated CO₂

Acknowledgments

Rob Jackson

Department of Biology and Nicholas School of the Environment, Duke University

Andrew McElrone

Post Doctoral Research Associate, Department of Biology, Duke University

Tracey Crocker

Lab Technician, Duke University

GCEP

Jeff Gaffney, Milton Constantin, Mary Kinney